



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

354. Proposed by G. PAASWELL, New York City.

The acceleration of an electric train is constant and equal to a feet per sec. per sec. Its braking or deceleration is variable and equal to the square root of the velocity. If the distance between stations is 5,000 feet, show that the acceleration must cease and braking ensue when the train is about 960 feet from the stopping point; also that the maximum velocity attained for a minimum time run is 88 m.p.h. and the time of run 54 seconds.

355. Proposed by HORACE OLSON, Chicago, Illinois.

A solid spheroid, axes a , a , b , is placed with its axis of revolution vertical. From its highest point, a particle is projected horizontally with a speed s . Where will it leave the spheroid, assuming that it slides on the surface without friction?

NUMBER THEORY.

272. Proposed by C. C. YEN, Tangshan, North China.

How many integers prime to n are there in each of the sets:

- (a) $1 \cdot 2, 2 \cdot 3, 3 \cdot 4, \dots, n(n+1);$
 (b) $1 \cdot 2 \cdot 3, 2 \cdot 3 \cdot 4, 3 \cdot 4 \cdot 5, \dots, n(n+1)(n+2);$
 (c) $\frac{1 \cdot 2}{2}, \frac{2 \cdot 3}{2}, \frac{3 \cdot 4}{2}, \dots, \frac{n(n+1)}{2};$
 (d) $\frac{1 \cdot 2 \cdot 3}{6}, \frac{2 \cdot 3 \cdot 4}{6}, \frac{3 \cdot 4 \cdot 5}{6}, \dots, \frac{n(n+1)(n+2)}{6}?$

273. Proposed by V. M. SPUNAR, Chicago, Illinois.

The ratio of the chances that all numbers ending in 1 or 9 and those ending 3 or 7 are composite is 3 : 2.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

477. Proposed by J. L. RILEY, Junior College, Stephenville, Texas.

Evaluate the product $(1 + r + r^2 + r^3)(1 + r^2 + r^4 + r^6) \dots (1 + r^{2^{n-1}} + r^{2^n} + r^{3 \cdot 2^{n-1}}).$

SOLUTION BY LOUIS O'SHAUGHNESSY, University of Pennsylvania.

Since

$$1 + r + r^2 + r^3 = \frac{1 - r^4}{1 - r}, \quad 1 + r^2 + r^4 + r^6 = \frac{1 - r^8}{1 - r^2}, \quad 1 + r^4 + r^8 + r^{12} = \frac{1 - r^{16}}{1 - r^4}, \quad \dots,$$

$$1 + r^{2^{n-1}} + r^{2^n} + r^{3 \cdot 2^{n-1}} = \frac{1 - r^{2^{n+1}}}{1 - r^{2^{n-1}}},$$

the required product is equal to the product of the following fractions:

$$\frac{1 - r^4}{1 - r} \cdot \frac{1 - r^8}{1 - r^2} \cdot \frac{1 - r^{16}}{1 - r^4} \cdot \frac{1 - r^{32}}{1 - r^8} \dots \frac{1 - r^{2^{n+1}}}{1 - r^{2^{n-3}}} \cdot \frac{1 - r^{2^n}}{1 - r^{2^{n-2}}} \cdot \frac{1 - r^{2^{n+1}}}{1 - r^{2^{n-1}}}.$$

The result, manifestly, is

$$\frac{(1 - r^{2^n})(1 - r^{2^{n+1}})}{(1 - r)(1 - r^2)}.$$

Also solved by H. C. FEEMSTER, PAUL CAPRON, ELIJAH SWIFT, and HORACE OLSON.